

# Earth's Trojan asteroid

Martin Connors<sup>1,2</sup>, Paul Wiegert<sup>3</sup> & Christian Veillet<sup>4</sup>


It was realized in 1772 that small bodies can stably share the same orbit as a planet if they remain near 'triangular points' 60° ahead of or behind it in the orbit<sup>1</sup>. Such 'Trojan asteroids' have been found co-orbiting with Jupiter<sup>2</sup>, Mars<sup>3</sup> and Neptune<sup>4</sup>. They have not hitherto been found associated with Earth, where the viewing geometry poses difficulties for their detection<sup>5</sup>, although other kinds of co-orbital asteroid (horseshoe orbiters<sup>6</sup> and quasi-satellites<sup>7</sup>) have been observed<sup>8</sup>. Here we report an archival search of infrared data for possible Earth Trojans, producing the candidate 2010 TK<sub>7</sub>. We subsequently made optical observations which established that 2010 TK<sub>7</sub> is a Trojan companion of Earth, librating around the leading Lagrange triangular point, L<sub>4</sub>. Its orbit is stable over at least ten thousand years.

The existence of Trojan asteroids of other planets raises the question of whether such companions could exist for Earth. Despite studies showing that such bodies could be relatively stable<sup>9</sup> and may wander relatively far from the Lagrange points<sup>5</sup>, they would dwell mostly in the daylight sky as seen from Earth, making detection difficult. Indeed, they hitherto have not been observed<sup>10,11</sup>. The launch of the Wide-field Infrared Survey Explorer (WISE) by NASA in 2009<sup>12</sup> provided improved viewing circumstances that made possible new detections of over 500 near-Earth objects<sup>13</sup>. WISE searched large areas of sky always 90° from the Sun, with high efficiency for asteroidal bodies and good astrometric accuracy. Examining WISE discoveries in the expectation that Earth co-orbital objects, possibly including a Trojan, could be found, resulted in two promising candidates, 2010 SO<sub>16</sub> and 2010 TK<sub>7</sub>. Both are larger than most co-orbital objects, being several hundred metres in diameter, and 2010 SO<sub>16</sub> is a horseshoe orbiter<sup>14</sup>. We identified 2010 TK<sub>7</sub> as probably being an Earth Trojan, on the basis of positions measured over a six-day arc in late 2010. Observations made at the University of Hawaii (D. Tholen, personal communication) and the Canada–France–Hawaii Telescope<sup>15</sup> in April 2011, after the object had for months been in an unfavourable position as seen from Earth, so greatly improved the knowledge of its orbit that we can state with certainty that 2010 TK<sub>7</sub> is an Earth Trojan.

The 'tadpole' motion of 2010 TK<sub>7</sub>, which is characteristic of Trojan asteroids, is shown in Fig. 1 in the frame co-rotating with Earth (see Supplementary Information for three-dimensional depictions of the motion). The 1-yr-averaged curve shows the centre of motion librating about L<sub>4</sub>, the Lagrange point 60° ahead of Earth. The period of this motion is at present 395 yr. Superposed on this is an annual motion or epicycle<sup>2,16,17</sup> (not shown for clarity). This mode of display emphasizes the longitudinal motion despite the enhanced radial scale: the asteroid's mean position drifts along the red line, from the 'head' of the tadpole, near Earth, to the far 'tail', where it is nearly on the opposite side of the Sun from the Earth. The relatively large eccentricity, of  $e = 0.191$ , results in an annual heliocentric radial motion between roughly 0.81 and 1.19 AU. The inclination of 2010 TK<sub>7</sub> is about  $i = 20.9^\circ$ , so there is significant motion perpendicular to Earth's orbital plane. The asteroid's eccentricity and inclination produce a large epicycle, which is responsible for the visibility of the object at the solar elongation of 90°, as observed by WISE; and it is now at the near-Earth end of the tadpole. In the present epoch, the longitude remains in the sector of L<sub>4</sub>, trapped

between Earth and L<sub>3</sub>. Interaction with Earth at the near-Earth end of the tadpole results in a rapid decrease in the object's semimajor axis,  $a$ , making it increase its angular speed (Kepler's third law) and outpace Earth. This is currently taking place. Slow resonant interaction at the other parts of the tadpole increases  $a$ , making the object slow gradually such that it again approaches Earth. In the current cycle, this will take place in the years AD 2050–2350, approximately. Repetition of this cycle leads to a sawtooth pattern in  $a$  as a function of time (Fig. 1c).

The present motion of 2010 TK<sub>7</sub> is well established, but there are inherent limits on our ability to compute orbits into the past or future. Chaos limits the accuracy of computations of this asteroid's position over timescales<sup>18</sup> greater than about 250 yr. However, we can still discuss the basic nature of its orbit with confidence by computing the motion of many 'dynamical clones' whose orbital parameters vary<sup>7</sup> within the limits set by observations. Approximately 1,800 yr in the past, and more than 5,000 yr in the future, the 100 clone orbits we




**Figure 1 | Orbital parameters of asteroid 2010 TK<sub>7</sub>.** **a**, Path over one Trojan libration from AD 2010 to 2405 in the co-rotating reference frame. In this frame, Earth is stationary and the average position of the asteroid librates about L<sub>4</sub> in a 'tadpole' orbit. Both Earth and the asteroid revolve about the Sun with periods close to 1 yr, and slow changes in their relative positions are best seen in the co-rotating frame. The difference between the asteroid's semimajor axis,  $a$ , and a circle of radius 1 AU (an astronomical unit (AU) is the Earth–Sun distance) is multiplied by a factor of 20 for clarity, and Earth and the Sun are not shown to scale. Black lines indicate  $a$  and longitude relative to Earth daily; the red curve shows the annual average. **b**, Longitude relative to Earth,  $\theta - \theta_E$ , over the period 420 BC to AD 4200. A 'jump' from L<sub>5</sub> libration to the present L<sub>4</sub> libration took place in around AD 400. Black and red lines indicate daily and averaged values, as in **a**. The grey band is the period of the present libration. **c**, Semimajor axis daily values. Initial conditions (best orbital solution) are given in Table 1.

<sup>1</sup>Athabasca University, 1 University Drive, Athabasca, Alberta T9S 3A3, Canada. <sup>2</sup>Department of Earth and Space Sciences, UCLA, Los Angeles, California 90095, USA. <sup>3</sup>Department of Physics and Astronomy, The University of Western Ontario, London, Ontario N6A 3K7, Canada. <sup>4</sup>Canada–France–Hawaii Telescope, Kamuela, Hawaii 96743, USA.

computed diverged sufficiently that we must say that the asteroid's precise behaviour cannot be predicted with certainty outside that  $\sim 7,000$ -yr span. The range of behaviour shown by the clones and, thus, possible for the real object includes making transitions to horseshoe modes and 'jumping' between Lagrange points. Short-term unstable libration about  $L_3$ , the Lagrange point on the other side of the Sun from Earth, can occur as a result of the asteroid's large inclination. Such orbits were theorized as early as 1920<sup>17</sup>, but no real object had until now been suspected to enter them.

Jumping from one Lagrange point to the other is a behaviour previously attributed to the Jupiter Trojan 1868 Thersites<sup>19</sup>, and was found in about half of the clone orbits. Here, the large eccentricity leads to longitudinal excursions (Kepler's second law), including when near  $L_3$ . In Fig. 2, these are shown to have allowed (in about AD 500) a rapid transition of 2010 TK<sub>7</sub> from  $L_5$  to the present  $L_4$  libration. The libration now remains only in the sector of  $L_4$  and is relatively stable, in a classic<sup>16</sup> Trojan pattern, although of large amplitude.



**Figure 2 | Semimajor axis versus relative longitude for 2010 TK<sub>7</sub>.**  
**a**, Libration during the period AD 1–800, featuring a 'jump' from libration initially about  $L_5$  (right) to the present libration around  $L_4$ . As in Fig. 1, black lines indicate daily values and red lines indicate the annual averages. When the asteroid is near a relative longitude,  $\theta - \theta_E$ , of about  $180^\circ$ , the annual excursions in relative longitude can cause it to cross  $L_3$ . This crossing can trigger a rapid transition or 'jump' between librational modes. Clone studies show that the chaotic behaviour of the asteroid is due mainly to a great sensitivity to non-resonant perturbations when near  $L_3$ . Libration about  $L_5$  results in an average longitude  $120^\circ$  different from that for libration about  $L_4$ . Such a large change resulting from small perturbations (when near  $L_3$ ) is characteristic of chaos.  
**b**, Present (AD 2010–2405) libration about  $L_4$ . The location of  $L_3$  is shown for reference but the relative longitude in the era after AD 800 does not cross it, resulting in the current stability of the orbit. The apparent banding in both panels is due to changes in semimajor axis and has a predominant period of roughly 12 yr; therefore, it is probably mainly caused by Jupiter perturbations.

**Table 1 | Heliocentric orbital elements of 2010 TK<sub>7</sub>**

Epoch	JD 2455600.5
Semimajor axis, $a$	1.0004078 AU
Eccentricity, $e$	0.1908177
Inclination, $i$	$20.87984^\circ$
Argument of perihelion	$45.86009^\circ$
Longitude of ascending node	$96.54190^\circ$
Mean anomaly	$20.30069^\circ$

Results in the figures were obtained using these initial conditions in the Mercury integrator<sup>24</sup> (verified in the near-present using the JPL Horizons system<sup>25</sup>). The RADAU<sup>26</sup> option was used with 1-d spacing for the eight planets, Pluto and the Earth–Moon barycentre. Clone studies included the eight planets and the Moon<sup>27</sup>, with variations<sup>7</sup> among the orbital elements of the order of the last significant digit shown. The Julian date (JD) shown corresponds to 0:00 UT on 8 February 2011.

Chaotic effects have a large role in the behaviour of this asteroid. Its sensitivity to small influences when in the vicinity of  $L_3$  allows the range of outcomes seen among the clones. The overall Trojan behaviour is dictated by 1:1 orbital resonance with Earth, but non-resonant effects of Jupiter are 80 times stronger than those of Earth when Jupiter is at the same celestial longitude as  $L_3$ . Such influences, demonstrated by the 'banding' seen in Fig. 2, alter the asteroid's chaotic behaviour. Many clone orbits make repeated transitions between the Lagrange points, such that the chaos can be stable<sup>20</sup>, with  $L_4$  and  $L_5$  each defining permitted regions of phase space. Knowledge of the orbit will improve as it is observed over the years, but its chaotic nature dictates that dynamics-based discussions of the origin and fate of 2010 TK<sub>7</sub>, and its relationship to other bodies, will necessarily remain statistical in nature.

Earth Trojan asteroids have been proposed as natural candidates for spacecraft rendezvous missions<sup>21</sup>. However, the large inclination of 2010 TK<sub>7</sub> results in a  $\Delta v$  of  $9.4 \text{ km s}^{-1}$  being required, whereas other near-Earth asteroids have values of  $\Delta v$  less than  $4 \text{ km s}^{-1}$ . The reported absolute magnitude, 20.7 mag, puts the diameter of 2010 TK<sub>7</sub> at 300 m with an assumed albedo of 0.1 (ref. 22), which makes it relatively large among the near-Earth asteroid population. No spectral or colour information is as yet available to determine whether the asteroid is in any other way unusual.

Received 11 April; accepted 27 May 2011.

- Wilson, C. in *Planetary Astronomy from the Renaissance to the Rise of Astrophysics. Part B: The Eighteenth and Nineteenth Centuries* (eds Taton, R. & Wilson, C.) 108–130 (Cambridge, 1995).
- Stacey, R. G. & Connors, M. A centenary survey of orbits of co-orbitals of Jupiter. *Planet. Space Sci.* **56**, 358–367 (2008).
- Connors, M. *et al.* A survey of orbits of co-orbitals of Mars. *Planet. Space Sci.* **53**, 617–624 (2005).
- Almeida, A. J. C., Peixinho, N. & Correia, A. C. M. Neptune Trojans and Plutinos: colors, sizes, dynamics, and their possible collisions. *Astron. Astrophys.* **508**, 1021–1030 (2009).
- Wiegert, P., Innanen, K. A. & Mikkola, S. Earth Trojan asteroids: a study in support of observational searches. *Icarus* **145**, 33–43 (2000).
- Wiegert, P., Innanen, K. A. & Mikkola, S. An asteroidal companion to the Earth. *Nature* **387**, 685–686 (1997).
- Brasser, R. *et al.* Transient co-orbital asteroids. *Icarus* **171**, 102–109 (2004).
- Connors, M. *et al.* Discovery of an asteroid and quasi-satellite in an Earth-like horseshoe orbit. *Meteor. Planet. Sci.* **37**, 1435–1441 (2002).
- Mikkola, S. & Innanen, K. A. Studies in solar system dynamics. II. The stability of Earth's Trojans. *Astron. J.* **100**, 290–293 (1990).
- Whitely, R. J. & Tholen, D. J. A CCD search for Lagrangian asteroids of the Earth–Sun system. *Icarus* **136**, 154–167 (1998).
- Connors, M. *et al.* Initial results of a survey of Earth's  $L_4$  point for possible Earth Trojan asteroids. *Bull. Am. Astron. Soc.* **32**, 1019 (2000).
- Wright, E. L. *et al.* The Wide-field Infrared Survey Explorer (WISE): mission description and initial on-orbit performance. *Astron. J.* **140**, 1868–1881 (2010).
- Mainzer, A. *et al.* Preliminary results from NEOWISE: an enhancement to the Wide-Field Infrared Survey Explorer for solar system science. *Astrophys. J.* **731**, 53 (2011).
- Christou, A. A. & Asher, D. J. A long-lived horseshoe companion to the Earth. *Mon. Not. R. Astron. Soc.* advance online publication, doi:10.1111/j.1365-2966.2011.18595.x (12 April 2011).
- Boulaide, O. *et al.* Development of MegaCam, the next-generation wide-field imaging camera for the 3.6-m Canada-France-Hawaii Telescope. *Proc. SPIE* **2008**, 657–668 (2000).
- Murray, C. D. & Dermott, S. F. *Solar System Dynamics* 63–110 (Cambridge Univ. Press, 1999).
- Moulton, F. R. *Periodic Orbits* 299–324, 151–198 (Carnegie Institute, 1920).
- Whipple, A. L. Lyapunov times of the inner asteroids. *Icarus* **115**, 347–353 (1995).

19. Tsiganis, K., Dvorak, R. & Pilat-Lohinger, E. Thersites: a 'jumping' Trojan? *Astron. Astrophys.* **354**, 1091–1100 (2000).
20. Milani, A. & Nobili, A. M. An example of stable chaos in the Solar System. *Nature* **357**, 569–571 (1992).
21. Stacey, R. G. & Connors, M. Delta-v requirements for earth co-orbital rendezvous missions. *Planet. Space Sci.* **57**, 822–829 (2009).
22. Bowell, E. *et al.* in *Asteroids II* (eds Binzel, R., Gehrels, T. & Matthews, M.) 524–556 (Univ. Arizona Press, 1989).
23. MPEC 2011-G37: Daily Orbit Update (2011 Apr. 5 UT). *IAU Minor Planet Center* (<http://www.minorplanetcenter.org/mpec/K11/K11G37.html>) (2011).
24. Chambers, J. E. A hybrid symplectic integrator that permits close encounters between massive bodies. *Mon. Not. R. Astron. Soc.* **304**, 793–799 (1999).
25. Giogini, J. D. *et al.* JPL's online solar system data service. *Bull. Am. Astron. Soc.* **28**, 1158 (1996).
26. Everhart, E. in *Dynamics of Comets: Their Origin and Evolution* (eds Carusi, A. & Valsecchi, G. B.) 185–202 (Kluwer, 1985).
27. Standish, E. M. *Planetary and Lunar Ephemerides DE406/LE406*. Tech. Report IOM 312.F-98-048 (NASA Jet Propulsion Laboratory, 1998).

**Supplementary Information** is linked to the online version of the paper at [www.nature.com/nature](http://www.nature.com/nature).

**Acknowledgements** We thank the WISE team and JPL and NEODyS (University of Pisa) data services. Support came from Canada's Natural Sciences and Engineering Research Council and Research Chairs. Problems with some 2010 TK<sub>7</sub> positions reported in ref. 23 were pointed out by T. Spahr and D. Tholen. We are grateful to them for data reductions provided, and to D. Tholen, M. Micheli and G. T. Elliot for observations made in support of this study.

**Author Contributions** The authors contributed equally to this work. M.C. and P.W. concentrated on dynamical calculations, and C.V. concentrated on observations and associated data reduction.

**Author Information** Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at [www.nature.com/nature](http://www.nature.com/nature). Correspondence and requests for materials should be addressed to M.C. ([martinc@athabascau.ca](mailto:martinc@athabascau.ca)).